

Amendments to the Specification:

Please replace the Brief Description of Figure 1, 2 and 6 on page 17 with the following descriptions.

E¹
-- Figure 1 Represents the primary structure of the peptides according to the invention
Shows the nucleic acid and amino acid sequences of the invention. Figure 1A-1 and 1A-2 show
the nucleic acid and amino acid sequence of SEQ ID Nos 1 and 4, respectively. Figure 1B-1,
1B-2, and 1B-3 show the nucleic acid and amino acid sequence of SEQ ID Nos 2 and 5,
respectively. Figure 1D-1 to 1D-3 show the nucleic acid and amino acid sequence of SEQ ID
Nos. 3 and 6, respectively.—

-- Figure 2 represents the amino acids sequence of the active human CCR5
chemokine receptor (SEQ ID NO: 5) according to the invention aligned with that of the human
CCR1 (SEQ ID NO: 9), CCR2b (SEQ ID NO: 7), CCR3 (SEQ ID NO: 8), and CCR4 (SEQ ID
NO: 10) receptors. Amino acids identical with the active CCR5 sequence are boxed.]--

E²
-- Figure 6 represents the structure of the mutant form of human CCR5 receptor.
Figure 6B shows the wild type amino acid sequence (SEQ ID NO: 11), and the location of the 32
base deletion mutation in the nucleic acid (SEQ ID NO: 12) and amino acid sequences (SEQ ID
NO: 13).]—

On page 30, please replace paragraph 2, extending from lines 5-32 on page 30 and lines
1-28 on page 31, with the following replacement paragraph;

E³
-- It is known that some individuals remain uninfected despite repeated exposure to HIV-1
[55, 56, 51]. A proportion of these exposed-uninfected individuals results from the
relatively low risk of contamination after a single contact with the virus, but it has been
postulated that truly resistant individuals do exist. In fact, CD4 lymphocytes isolated
from exposed-uninfected individuals are highly resistant to infection by primary M-
tropic, but not T-tropic HIV-1 strains. Also, peripheral blood mononuclear cells (PBMC)
from different donors are not infected equally with various HIV-1 strains [57-59]. Given
the key role played by CCR5 in the fusion event that mediates infection by M-tropic
viruses, it is postulated that variants of CCR5 could be responsible for the relative or
absolute resistance to HIV-1 infection exhibited by some individuals, and possibly for the
variability of disease progression in infected patients [66]. The Inventors selected three
HIV-1 infected patients known to be slow progressors, and four seronegative individuals
as controls; the full coding region of their CCR5 gene was amplified by PCR and

sequenced. Unexpectedly, one of the slow progressors, but also two of the uninfected controls, exhibited heterozygosity at the CCR5 locus for a biallelic polymorphism. The frequent allele corresponded to the published CCR5 sequence, while the minor one displayed a 32 bp deletion within the coding sequence, in a region corresponding to the second extracellular loop of the receptor (Fig. 6). The figure 6 is the structure of the mutant form of human CC-chemokine receptor 5. a. The amino acid sequence of the non-functional Δ ccr5 protein is represented. The transmembrane organisation is given by analogy with the predicted transmembrane structure of the wild-type CCR5. Amino acids represented in black correspond to unnatural residues resulting from the frame shift caused by the deletion. The mutant protein lacks the last three transmembrane segments of CCR5, as well as the regions involved in G protein-coupling b. Nucleotide sequence of the CCR5 gene surrounding the deleted region, and translation into the normal receptor (top) or the truncated mutant (ccr5, bottom). The 10-bp direct repeat is represented in italics. The full size coding region of the CCR5 gene was amplified by PCR, using 5'-TCGAGGATCCAAGATGGATTATCAAGT-3' (SEQ ID NO: 14) and 5'-CTGATCTAGAGCCATGTGCACAACTCT-3' (SEQ ID NO: 15) as forward and reverse primers' respectively. The PCR products were sequenced on both strands using the same oligonucleotides as primers, as well as internal primers, and fluorochrome-labelled didoxynucleotides as terminators. The sequencing products were run on an Applied Biosystem sequencer, and ambiguous positions were searched along the coding sequence. When the presence of a deletion was suspected from direct sequencing, the PCR products were cloned after restriction with *Bam*HI and *Xba*I endonucleases into pcdna3. Several clones were sequenced to confirm the deletion. The deletion was identical in three unrelated individuals investigated by sequencing-

On page 33, please replace paragraph 3 extending from lines 24-32 on page 33 and lines 1-19 on page 34 with the following replacement paragraph;

The figure 8 represents genotyping of individuals by PCR and segregation of the CCR5 alleles in CEPH families. a, Autoradiography illustrating the pattern resulting from PCR amplification and *Eco*RI cleavage for individuals homozygous for the wild-type CCR5 allele (CCR5/CCR5), the null Δ ccr5 allele (Δ ccr5/ Δ ccr5), and for heterozygotes (CCR5/ Δ ccr5). A 735 bp PCR product is cleaved into a common band of 332 bp for both alleles, and into 403 and 371 bp bands for the wild-type and mutant alleles, respectively. b, Segregation of the CCR5 alleles in two informative families of the CEPH. Half-black and white symbols represent heterozygotes and wild-type homozygotes, respectively. For a few individuals in the pedigrees, DNA was not available (ND: not determined). PCRs were performed on genomic DNA samples, using 5'-CCTGGCTGTCGTCCATGCTG-3' (SEQ ID NO: 16) and 5'-CTGATCTAGAGCCATGTGCACAACTCT-3' (SEQ ID NO: 17) as forward and reverse primers respectively. Reaction mixtures consisted in 30 μ l of 10 Mm Tris-Hcl buffer Ph 8.0, containing 50 Mm Kcl, 0.75 Mm MgCl₂, 0.2 Mm dCTP, dGTP and dTTP, 0.1 Mm dATP, 0.5 μ i [a-³²P]-DATP, 0.01% gelatine, 5% DMSO, 200 ng target DNA, 60 ng of each of the primers and 1.5 U Taq polymerase. PCR conditions were: 93 °C for 2 min 30; 93 °C for 1 min, 60 °C for 1 min, 72 °C for 1 min, 30 cycles; 72 °C for 6 min. After the PCR reaction, the samples were incubated for 60 min at 37 °C with 10 U

E⁴

*Eco*RI, and 2 μ l of the denatured reaction mixture was applied onto a denaturing 5% polyacrylamide gel containing 35% formamide and 5.6 M urea. Bands were detected by autoradiography.
